

Fertimix-Fe

CORRECTOR OF LACKS OF FER QUELATE (LIQUID)

CONTENTS GUARANTEED: (From gross matter)

Iron (Fe)7 % w/w(9,8 %w/v)

Density.....1'4 gr/cc

PH.....2'1

DEFINITION

The microelements and other microelements are necessary for plants life, even when we can find in small quantities.

The function of these elements is took part of the cellular mass and improve metabolic reactions.

Fertimix-Fe is a liquid formulation with acid heptagluconic and EDDHA for the preventive and curative control of the chlorosis.

The principal functions of the iron in the plant are the formation of the chlorophyl, proteins, fixing of the nitrogen and it works in the breathing.

PREPARATION OF MIXTURES AND COMPATIBILITIES

Before preparing a final mixture is suitable make some compatibility test. In case of doubt contact with the Technical Department. **Fertimix-Fe** has compatibility with the most of fertiliser N-P-K and phythosanitarries of normal usage. Avoid mixtures with strong acids neither products of reaction very alkalinised.

APPLICATIONS

Fertimix-Fe can be applied in leaves or in soils, the range of stability of the product is between a pH from 3.7 to 10.2. As sensitive cultures to the application of the iron are fruit trees, horticulture and vine tree.

Recommended doses can change in function of the application:

Leaves application: 6-12 l/Ha distributed in the cycle of the culture.

Soil application: 8-20 l/Ha distributed in the cycle of the culture.

NOTES AND ADVISES

The recommendations and information that we give are the result of wide and severe test. However, in the utilisation could take part a lot of factors which escape to our control (preparation of mixtures, application, weather, etc.)The company assures the composition, formulation and content. The user will be responsible of the damage caused (fault of efficacy, toxicity in general, residues, etc., for not to follow the instructions in the label.

FERTILIZER C.E.

Contact with our technical service if you have any doubt

SUGGESTIONS FOR USE

CULTURE	DOSES		WAY AND TIME OF APPLICATION
	Leaves	Soil	
Fruit trees	150-200 cc/100 l/water	2-3 L/Ha / application	Make from 3 to 4 applications from the begining of the sprout time.
Horticulture and Strawberry	200-300 cc/100 l/water	2.5-3 L/Ha /application	Make from 4 to 5 applications since the transplant.
Hidroponic	150-250cc/100 l/water	2-3 L/m ³ of solution mother concentrated 100 times.	Make 1 application every week since one month before the harvesting.
Vine Trees	150-250 cc/100 l/water	2-2.5 L/Ha /application	Make from 2 to 3 applications after the beginning of the sprout time.
Citrus	200-250 cc/100 l/water	2-3 L/Ha/application	Make from 2 to 3 treatments distributed in springtime and sprout time in summer.
Ornamentals	50-100 cc/100 l/water	1.5-2 L/Ha /application	Make from 2 to 3 applications making them before the flowerings.
Plane trees	200-250 cc/100 l/water	3-5 L/Ha /application	Make from 2 to 4 applications distributed at the end of winter and at the end of summer.
Other kind of crops	150-300 cc/100 l/water	2-3 L/Ha /application	Make from 2 to 3 applications since the beginning of the culture.