

Granfol-K

POTASSIUM PHOSPHYTE

CONTENTS GUARANTEED: (From gross matter)

Phosphorous (P₂O₅)..... 32 % w/w(45,76 % w/v)
Potassium(K₂O)..... 22 % w/w (31,46 % w/v)
Density.....1'43 gr/cc
Ph.....5

DEFINITION

Granfol-K is a liquid formulation of Phosphorous and potassium. The presence of Potassium allows a bigger fattening of fruits. The Ion Phosphyte allows the transformation of PO₃⁻³ in PO₄⁻³, creating in the plants metabolism some natural protection, so it is advisable its employ in crops affected by fungus endoparasitaries as Oomicets (fungus producers of watered in citrus, rottenness of wooden crops, mildius of vines and horticultural, etc.)

APPLICATIONS

Granfol K is ready for leaves application and for the soil application, applying in cultivation of citric, fruit trees, tropical fruit, vine, in horticulture : potato, tomato and strawberry.

Recommended doses can change in function of the application:

Leaves application: 6-8 l/Ha /year

Soil application: 6-12 l/Ha/ year

FERTILIZER C.E.

PREPARATION OF MIXTURES AND COMPATIBILITIES

Before preparing a final mixture is suitable make some compatibility test. In case of doubt contact with the Technical Department. **Granfol-K** has compatibility with the most of fertilisers an phythosanitarities of normal use. Don't mix with products highly alkalinised, mineral oils, *Dimetoato* or *Dicofol*, *Dinocap* and preparations with Cuprous.

NOTES AND ADVISES

The recommendations and information that we give are the result of wide and severe test. However, in the utilisation could take part a lot of factors which escape to our control (preparation of mixtures, application, weather, etc.)The company assures the composition, formulation and content. The user will be responsible of the damage caused (fault of efficacy, toxicity in general, residues, etc., for not to follow the instructions in the label.

Contact with our technical service if you have any doubt

SUGGESTIONS FOR USE

CULTURE	DOSES		WAY AND TIME OF APPLICATION
	Leaves	Soil	
Citric and Fruit trees	250-350 cc/100 l/water	4 -5 L/Ha and irrigation	Apply in spring time (March, April) and after harvest.
Vine and olive tree	250-300 cc/100 l/water	3-5 L/Ha and irrigation	Make from 2 to 3 applications from the sprout time.
Horticulture and strawberry	200-300 cc/100 l/water	6-7 L/Ha and irrigation	Make applications divided along the cycle of the crop.
Industrial: Potatoe, bet-root, tomato, etc	250-300 cc/100 l/water	2-3 L/Ha and irrigation	Make 2 applications when there is enough foliate mass.
Ornamentals	300 cc/100 l/water	5-6 L/Ha and irrigation	Make from 3 to 4 applications distributed in the cycle of the crop.
Other kind of crops	250 cc/100 l/water	6 L/Ha and irrigation	Make from 2 to 3 applications shared in the cycle of the crop.