

Hortumus

LIQUID HUMIC ACID

CONTENTS GUARANTEED: (From gross matter)

Total Humic extract.....	20,5 % w/w (24,6% w/v)
Humic Acids.....	10,5 % w/w(12,6% w/v)
Fulvic Acids.....	10 % w/w (12% w/v)
Density.....	1'2 gr/cc
PH.....	11

DEFINITION

Hortumus is an humic amendment that, for its rich composition in Humic Acids, is specially suitable for improve the structure of tired soils and very mineralised, helping the liberation of blocked nourishment and stimulating the capacity of retention of the soil. Likewise, its action of vegetative stimulation on the roots and in the aerial parts of the plants, allows a better balanced development that have repercussion on increasing productions.

APPLICATIONS

Hortumus can be applied by fertirrigation at 10 L/HAS. to the 15 days after transplanting and subsequently to ration weekly the remainder of the dose to apply.

Also can be utilized by leaves at 0,3% (0,3 Litters in 100 Litters of water). Is very recommendable to utilize it with systematic herbicides therefore power its action and aid to its.

PREPARATION OF MIXTURES AND COMPATIBILITIES

Before preparing a final mixture is suitable make some compatibility test. In case of doubt contact with the Technical Department. **Hortumus** has compatibility with the most of fertilisers and phythosanitariaries of normal use. Don't mix with products of acid reaction (sulphuric acid, nitric, phosphoric, etc.), as well as with products which contains Calcium or Magnesium.

NOTES AND ADVISES

The recommendations and information that we give are the result of wide and severe test. However, in the utilisation could take part a lot of factors which escape to our control (preparation of mixtures, application, weather, etc.)The company assures the composition, formulation and content. The user will be responsible of the damage caused (fault of efficacy, toxicity in general, residues, etc.), for not to follow the instructions in the label.

Contact with our technical service if you have any doubt

USES RECOMMENDED

CULTURE	DOSE BY SOIL	WAY AND TIME OF APPLICATION
Fruits and citrus trees	25-40 L/ Ha and year	To apply from sprouting to 1 month before ha
Grapes and vines	15-30 L/ Ha and year	To apply from sprouting.
Strawberries	20-30 L/ Ha and year	To apply during the crop cycle.
Banana	40-80 L/ Ha and year	To apply during the crop cycle.
Industrials	15-20 L/ Ha and year	To apply when culture has got from 6 to 8 lea
Ornamentals	20-40 L/ Ha and year	To apply during the crop cycle
Cultures in general	20-40 L/ Ha and year	To apply during the crop cycle



ARVENIS AGRO, S.A.
Carretera Castellón Km.212,1
50740 Fuentes de Ebro
ZARAGOZA (ESPAÑA)
Tel. 00 34 976 169 181
Fax 00 34 976 169 183
e-mail: mail@arvensis.com
www.arvensis.com