

Rhizum

SPECIAL PRODUCT CONTAINING AMINOACIDS

CONTENTS GUARANTEED: (From gross matter)

Free Amino acids.....	7,5 % w/w(8,4 %w/v)
Total nitrogen	5 % w/w(6' % w/v)
Ureic nitrogen	2,25 % w/w(2'7 % w/v)
Organic nitrogen	2,75 % w/w(3,2 % w/v)
Phosphorous (P ₂ O ₅).....	4% w/w (7,2% w/v)
Potassium (K ₂ O).....	4 % w/w (6,2% w/v)
Organic matter.....	30 % w/w(24 % w/v)
Density.....	1,2 gr/cc
pH.....	6,5

DEFINITION

Rhizum is a formulation which comes from vegetable extracts holding seaweed extract, humic-fulvic acid, polisacarids and organic matter, that reactive the root system by natural auxins.

The objective is to advance in taking root and in vegetable growth through a quick protean synthesis. As a result of that, cultivation advance fruit production.

APPLICATIONS

Rhizum is developed to be applied in any kind of irrigation system(dripping, sprinkling and gravity)

As a general recommendation this product must be applied 8-10 days after cultivation transplant in horticultural cultivations. In fruit cultivation, vine and nursery, it will be applied with the first plantation irrigation or with the first irrigation ant the sprout beginning.

PREPARATION OF MIXTURES AND COMPATIBILITIES

Before preparing a final mixture is suitable make some compatibility test. In case of doubt contact with the Technical Department. **RHIZUM** has compatibility with the most of NPK fertilisers of normal use. Don't mix with products of acid reaction (sulphuric acid, nitric, phosphoric, etc)

NOTES AND ADVISES

The recommendations and information that we give are the result of wide and severe test. However, in the utilisation could take part a lot of factors which escape to our control (preparation of mixtures, application, weather, etc.)The company assures the composition, formulation and content. The user will be responsible of the damage caused (fault of efficacy, toxicity in general, residues, etc.), for not to follow the instructions in the label.

Contact with our technical service if you have any doubt

SUGGESTIONS FOR USE

CULTURE	DOSES	WAY AND TIME OF APPLICATION
	Soils	
Citric and Fruit trees	4-5 l/Ha and application	Apply with first irrigation and a second application in the sprout time.
Horticulture	3-5 l/Ha and application	Make the first application before transplanting
Industrials	3 l/Ha and application	Make the first application when the culture has got two or three leaves, and 20 days before.
Banana trees	5 l/Ha and application	Make the first application at the end of winter
Ornamentals	5 l/Ha and application	Make from 2 to 3 applications shared in the cycle of the crop.
Vine and olive tree	4-5 l/Ha and application	Apply with first irrigation and a second application in the sprout time.
Tropicals	5 l/Ha and application	Make the first application at the end of winter.
Other kind of crops	4-6 l/Ha and application	Make from 2 to 3 applications shared in the cycle of the crop.